

MATERIALS AND METHODS

This was a survey conducted to understand the eating habits of primary school students from standards 4–6 within the age range of 10–12. Data collection from a large number of students was necessary to generalize the results. There were approximately 554, 994 primary school students (standards 1–6) in Selangor District (*Department of Statistics, 2015*). The number of students would be fewer (not more than 350,000) from standards 4 to 6. In accordance with Gay, if the population is 50,000, a sample size of 1% would be more than adequate (*Gay, 1996*). *Krejcie & Morgan (1970)* also determined a sample size of 381 for a population of 50,000, 382 for a population of 75,000 and 384 for a population of 100,000. They also suggested a sample size of 1% is adequate if the population is more than 100,000. Thus, the sample size (400 students) for this study is more than adequate to truly represent the total population as total students from standards 4–6 will not be more than 350,000 in Selangor District of Malaysia.

Participants

There are two types of public schools at the primary level: the National Schools and the National Type Schools. In the National Schools, the medium of instruction is the Malay language, which is the national language. The medium of instruction in the National Type Schools is either the Chinese language or the Tamil language. In both types of schools, the Malay language is a compulsory subject. English is compulsory and is taught as a second language in all schools. Chinese, Tamil and indigenous languages are also offered as subjects in the National Schools. All three types of public schools are included in this study; these comprise Sekolah Kebangsaan (National Schools), Sekolah Kebangsaan Jenis Tamil (National Type Schools) and Sekolah Kebangsaan Jenis Cina (National Type Schools). Invitations were sent to the respective schools until the number of schools required was achieved. In response to the invitations, the principals were given the responsibility to address the matter with the teachers and students involved in the study. After seeking agreement from the school to participate in the study, a number of teachers involved in the subject of *Pendidikan Jasmani dan Kesihatan (PJK)*, or Physical and Health Education, were asked to choose a class taught from either standards 4, 5 or 6 in order to obtain answers to the questionnaire.

Instrumentation: validity and reliability

A structured bilingual version of the questionnaire, comprising Malay and English, was used for the study. A student questionnaire was constructed for the purpose of assessing students in standards 4, 5 and 6 on the current knowledge, attitudes and practices about healthy eating.

The questionnaire was adapted from the Hong Kong Department of Health (*Department of Health, 2009*), and a pilot study was conducted to either modify or omit questions to suit the context of Malaysian schools. A pilot study was conducted in February 2015 to assess the content validity and comprehensibility of the measurement tools and to test the logistics of the study. Apart from that, before the study the opinions of experts have been taken and the questionnaire was corrected based on their suggestions

to improve the validity. A total of 45 students from three primary schools participated in the pilot study. Over 15 students were invited from each school with the selection of five students from each Grade. These three schools were excluded from the final study. The final questionnaire was revised based on the feedback from the pilot study. The Cronbach's Alpha of the instrument was between 0.60–0.90. This pilot study shows that majority of the questions are valid and produce reliable results for analysis; thus, no change was needed for the actual survey.

Procedure

As the primary school children are involved, permission has been taken from University of Malaya Ethics Committee. Between April 15, 2015 and April 19, 2015, 16 participating schools were identified, and the questionnaires were distributed to the students. The representatives involved in this study distributed the questionnaire to all the classes of standards 4, 5 and 6 that were chosen. All the students involved were asked to complete the questionnaire during a classroom session. The schools were given four weeks to conduct the questionnaires. The completed questionnaires were collected by the teachers. A representative performing the study collected all the questionnaires from all the schools involved. The completed questionnaires were collected from the schools between May 6, 2015 and June 8, 2015.

Data analysis

Descriptive statistics is used for all data representation of all survey questions. Generally, descriptive statistics is the easiest form of summarizing data in a presentable format. This simple statistics is performed in Statistical Package for Social Science version 20.0. Some of the most common function used for this survey is frequency, percentage, standard deviation and mean.

RESULTS

Rate of responses

A total of 600 questionnaires were distributed by the teachers to the students in standards 4, 5 and 6 at the 16 participating schools. A total number of 420 answered questionnaires were returned, resulting in a 70% response rate. Twenty of these questionnaires were unanswered, and the remaining 400 were used in the analysis. Respondents' demographics [Table 1](#) shows that the majority of respondents (40%) were students in standard 6. The respondents also included more male students (53.5%) and more students belonging to the Malay race (65.8%).

Students' knowledge of healthier food and their preferences

In identifying knowledge about healthy food, six pairs of food were given in the questionnaire. The students were asked to choose the healthiest food among the six pairs. [Table 2](#) indicates that the majority of students understood what healthy foods were based on the correct choices made.

Table 1 Characteristics of 400 children aged 10–12 years in standards 4–6 in the state of Selangor, Malaysia.

Variables	No. of students	Percentage
Grade (n = 400)		
Standard 4	143	35.8
Standard 5	97	24.3
Standard 6	160	40.0
Gender (n = 400)		
Male	214	53.5
Female	186	46.5
Race (n = 400)		
Malay	263	65.8
Chinese	44	11.0
Indian	93	23.3
School district (n = 400)		
Selangor	400	100

Table 2 Knowledge of healthier food and their preferences of 400 children aged 10–12 years in standards 4–6 in the state of Selangor, Malaysia.

Students' knowledge of healthier food choices				Students' food preferences			
Healthier choice	%	Unhealthier choice	%	Healthier choice	%	Unhealthier choice	%
Soya sauce drumsticks	55.0	Deep fried drumsticks	45.0	Soya sauce	48.7	Deep fried drumsticks	51.3
Pure orange juice	86.3	Carbonated drinks	13.7	Pure orange juice	79.3	Carbonated drinks	20.7
Raisin whole-meal	82.0	Hot dog	18.0	Raisin whole-meal	52.3	Hot dog	47.7
Yogurt	80.0	Ice cream	20.0	Yogurt	51.5	Ice cream	48.5
Spaghetti with fresh tomato sauce	77.3	Hamburger and fries	22.7	Spaghetti with fresh tomato sauce	39.8	Hamburger and fries	60.2
Fried noodles with vegetables	79.0	Fried noodles with chicken	21.0	Fried noodles with vegetables	43.8	Fried noodles with chicken	56.2

In identifying the students' food preferences, the same six pairs of food and drink as the knowledge questions were provided. Table 2 also shows that there was a difference in the answers between the students' preferences and their perceived opinions on what healthy foods are. The percentages of students' food preferences were generally lower. For example, although many students perceived that deep fried drumsticks, hamburgers and fries and fried noodles and chicken are unhealthy foods, more than half of the respondents still preferred eating those foods despite the fact that they are unhealthy choices.

Reasons for not consuming breakfast

A total of 67%, or 268 students, had consumed breakfast on the day of the survey. The students who did not consume breakfast showed different reasons. Table 3 shows that an alarming figure of 56.8% of the students did not have breakfast because they did not have enough time, and 22.7% were not in the habit of having breakfast. Another concerning reason was that nobody had prepared breakfast for 7.6% of them.

Table 3 Reasons for not having breakfast of 400 children aged 10–12 years in standards 4–6 in the state of Selangor, Malaysia.

Reason	No. of students	Percentage
I do not have time	75	56.8
I am on a diet	2	1.5
I want to save money	0	0.0
I am not used to having breakfast	30	22.7
I do not have the appetite to eat in the morning	15	11.4
No one has prepared breakfast for me in the morning	10	7.6
Other reasons	0	0.0

Table 4 Eating habits of various food categories on average every day of 400 children aged 10–12 years in standards 4–6 in the state of Selangor, Malaysia.

Food category	Eating frequency					
	No. of students	More than twice	Twice	Once	Never	Do not know
Fruits	400	47.0	18.8	26.0	5.8	2.4
Vegetables	400	44.5	24.0	21.0	6.8	3.7
Dairy products	400	30.3	18.5	31.8	14.8	4.6
Meat, chicken, fish	400	58.5	21.5	14.5	3.0	2.5
Grains	400	63.8	17.3	11.3	3.5	4.1
Fried and deep-fried food	400	37.0	23.8	28.8	7.3	3.1
Drinks with added sugar	400	15.3	20.3	40.0	20.5	3.9
Food high in sugar	400	20.2	16.2	40.1	16.0	7.5
Food high in salt	400	12.0	11.0	30.5	40.0	6.5
Food high in fat	400	24.5	17.5	39.8	11.8	6.4

Students' daily dietary habits

The students were required to evaluate their daily dietary habits during the week prior to answering the questionnaire. According to [Table 4](#), the percentage of students consuming fruits and vegetables more than twice was, on average, between 44.5 and 47.0%. However, there was a high occurrence of unhealthy eating habits, such as having deep-fried food and consuming foods high in sugar, salt and saturated fat. Between 60% and 92.7% of the students did not reach the recommended frequencies, as shown in [Table 5](#).

Students' knowledge of general healthy eating

Based on the two questions asked, there was a lack of healthy eating knowledge by the students because only 27.5% correctly chose grains as the type of food that should be provided most often in a healthy lunchbox and only 35.8% knew the correct recommended servings of fruits and vegetables ([Table 6](#)).

Students' snack eating habits and sources

Students were shown pictures of snacks before answering whether they had the habit of eating them. The majority of students (73.3%) had the habit of snacking; they mostly

Table 5 Percentage who reached the corresponding recommended frequency of 400 children aged 10–12 years in standards 4–6 in the state of Selangor, Malaysia.

Recommended eating frequency of various food categories	No. of students	No. of students/percentage of students	
		Reached recommended frequency	Not reached recommended frequency
Fruit \geq 2 times	400	263 (65.8%)	137 (34.2%)
Vegetables \geq 2 times	400	274 (68.5%)	126 (31.5%)
Dairy products \geq 1 time	400	322 (80.5%)	78 (19.5%)
Meat, chicken, fish \geq 1 time	400	320 (80.0%)	80 (20.0%)
Grains \geq 2 times	400	255 (63.8%)	145 (36.2%)
Not having fried and deep-fried food	400	29 (7.3%)	371 (92.7%)
Not having drinks with added sugar	400	82 (20.5%)	318 (79.5%)
Not having food high in sugar	400	63 (15.8%)	337 (84.2%)
Not having food high in salt	400	160 (40.0%)	240 (60.0%)
Not having food high in fat	400	47 (11.8%)	353 (88.2%)

Table 6 Percentage of students correctly answering the general healthy eating messages.

Healthy eating messages	No. of students	Percentage
A healthy lunchbox should have more grains	110	27.5
A person must consume two servings of fruits and three servings of vegetables daily	143	35.8

had homemade snacks (86.5%), while 2.3% bought snacks from the school co-operative shop. Others sources of snacks included outside shops and supermarkets (11.2%).

Matters for the students' consideration in selecting their food

In the event of choosing food, students were particular regarding the cleanliness of the food: 65.8% responded to cleanliness as a high priority, followed by 12.3% responding that the preference of parents determined food choice, as shown in [Table 7](#).

Students' perceived eating habits

[Figure 1](#) shows that a total of 45.5% of students viewed themselves as having very healthy eating habits, and only 1% considered their eating habits to be very unhealthy

Students' awareness of the Integrated School Health Program (ISHP)

More than 45% of the students had heard of the ISHP, whereas 55% did not know about the program.

Support in promoting healthy eating at schools

A total of 77.8% of the students stated that they supported healthy eating at school, whereas only 8.8% of the students were unsupportive of healthy eating. Another 13.4% did not know whether there was any program of promoting healthy eating at schools.

Table 7 Consideration for choosing food of 400 children aged 10–12 years in standards 4–6 in the state of Selangor, Malaysia.

Matters for consideration	No. of students	Percentage
Cleanliness	263	65.8
Taste	23	5.7
Good for health	17	4.2
Freshness	43	10.7
Price	4	1.0
Easy to get	1	0.3
Preference of parents	49	12.3
Choice of other schoolmates	0	0.0




Figure 1 Students' perception of their eating habits: data of 400 children aged 10–12 years in standards 4–6 in the state of Selangor, Malaysia.

Participation in school activities promoting healthy eating

Based on the students' responses regarding their participation in school activities that promoted healthy eating in the past year, only 13% of the students claimed to have been involved. Nearly half of the students (44.8%) claimed they were not sure of their participation in these types of activities, and 29.3% of the students reported that no such activities were ever organized in their schools.

DISCUSSION

Thus, the ISHP is a program that is still considered important in promoting a healthy eating environment. It is of utmost importance for students to learn healthy eating habits at an early age, thus defining the purpose of implementing the program at the primary level (*Gourdet et al., 2014*). This survey provides an opportunity to identify the dietary patterns of students and the elements that affect their eating habits. Findings show that the students understand the definition of healthy food but still 50% of the students consume deep-fried drumsticks and hamburgers containing a high amount of

saturated fat and cholesterol. Approximately 60% of the students have knowledge of the correct intake of foods such as vegetables, fruits, meats and grains. At the same time, a high percentage of students consume foods that are fried and contain sugar, salt and fat. Parents play an important role in determining the types of snacks as 86.5% of students responded that the snacks they consumed were taken from their homes. Another finding of students not having breakfast (67%) may cause serious health concern in later life.

Students' perceptions towards eating frequencies in various food categories

In terms of the students' practice in eating healthy foods, there was a high correct response rate regarding the recommendations of the eating frequency in the various food categories. The students responded well in terms of the correct intake of foods such as vegetables, fruits, meats and grains, with an average correct response rate of more than 60%. However, there were a high percentage of students who consumed foods that were fried and contain sugar, salt and saturated fat. This finding is in line with the high percentage of students preferring fried drumsticks in the question regarding the pairs of food. Based on these results, culture is an aspect that can be examined in terms of the types of food consumed. Malaysia is a country that has infused its food based on the different racial backgrounds. Most of the food that can be found and bought in Malaysia is prepared in ways that can be seen as unhealthy. This finding is supported by [Noraziah & Mohd Azlan \(2012\)](#), who stated that current Malaysian foods are sweet, oily and fatty, which have quality in taste but does not consider the aspect of health. Furthermore, with parents spending more time at work, students are exposed to foods that are prepared quickly, such as microwaved food in boxes and eating at fast food restaurants, where food can easily be ordered and sent directly to homes. As a result of these factors, the types of food consumed contain ingredients that are high in sugar, salt and fat, leading to unhealthy eating behaviors.

Problems students face in consuming breakfast

Breakfast is considered the most important meal of the day because it helps an individual to work and think during the early part of the day ([Pandey et al., 2013](#)). Without breakfast, students may not have the energy to think and perform tasks in class. According to the survey, only 67% of students had breakfast on the day of the survey. For those who did not have breakfast, 56.8% responded that not having enough time was their reason for missing breakfast. As such, [Singleton & Rhoads \(1982\)](#) also identified a lack of time (43%) as the main factor for not having breakfast. This effect may be due to the problem of time management. In most school systems, school sessions begin as early as 7:30 in the morning, thus requiring students to be in school before 7:30. Students must wake up early and prepare themselves for school. Students using transportation, such as school buses, must prepare themselves even earlier because the buses run early in the morning to pick up all the students in different areas to bring to school. As a result, some students have no time for breakfast and may wait until the morning recess to eat.

Students' perceptions and action towards healthy food

According to the survey, the majority of students were able to correctly identify the six pairs of food with each pair having a minimum of 55% correct answers. This finding indicates that the students understood the definition of healthy food and the types of foods that are considered healthy. However, in a question involving those same food pairs and the students' choice of food, there was a difference in the percentage of correct choices. Although the students knew that foods such as deep-fried drumsticks and hamburgers contain a high amount of saturated fat and cholesterol, these food are still consumed by the students, with more than 50% of the survey population reporting consumption of these foods. Only having knowledge about healthy eating habits is not sufficient; the practical aspect should also be emphasized to students, because there is no purpose of learning without implementing what has been taught (*Woodruff & Hanning, 2009*).

Students' habit of snacking

Snacks can be classified as foods that are eaten between meals or consumed as light meals and can be either healthy or unhealthy. Healthy snacks, such as fruits, vegetables and plain water, contribute to healthy eating and contain no saturated fat or artificial flavors. However, snacks such as chips and sweets contain an excess of ingredients, such as salt and sugar, that can lead to obesity. Moreover, several studies have indicated that the consumption of unhealthy snacks is one of the main contributors to being overweight (*de Graaf, 2006*). It was found that 73.3% of students have the habit of snacking, and 86.5% of students responded that the snacks they consumed were taken from their homes. This finding means that parents play an important role in determining the types of snacks that their children are eating. Parents should set an example for their children and differentiate between healthy and unhealthy food. This conclusion is further supported by *Noraziah & Mohd Azlan (2012)* who stated that families who are poor role models influence the perception of the types of food that their children eat.

Students' factors in choosing food

In choosing the food that they eat, two major factors contributed to the students' decision: cleanliness (65.8%) and the preference of their parents (12.3%). Cleanliness not only refers to the food but also to the way the food was prepared and the cleanliness of the premises where the food was bought. This finding shows that students are aware of the importance of cleanliness because contaminated food may cause food poisoning, cases of which have been reported in schools. As stated earlier, parents play an important role in determining the types of food their children eat. The results show that parents are aware of their children's eating habits and have influence on the types of food that their children eat. With the influence of parents, children have better guidance in the foods that they consume, which may prevent the consumption of unhealthy foods (*Cheng et al., 2008*).

Students' perceptions of the ISHP and the implementation of health programs

According to the survey, only 45% of the students had heard of the ISHP that was implemented by the Ministry of Education. Furthermore, 28.2% of students had never heard of the program. This result was considered alarming because 40% of the respondents were in standard 6 and will soon graduate from primary school. Most of the students have been in primary school for five or six years but had never heard of the program, which may be due to a failure of the school administrators to clarify this program to the students. It is important that the students are always briefed and reminded to help them understand the importance of health and to incorporate awareness into the students' learning process. Failing to implement the program in schools deprives students of knowledge related to health issues. As such, *Basch (2010)* and *Egner, Oza-Frank & Cunningham (2014)* stated that in accordance to scientific reviews, schools that conduct health programs can create positive effects on students' educational outcomes and health outcomes. However, 77.8% of the students supported their school in events promoting healthy eating, which meant that the students had a positive attitude towards healthy eating events and showed an interest in participating in those activities. Nevertheless, due to a lack of clarity, 44.8% of the students were uncertain if they had joined the program in the previous year, and 29.2% stated that their schools had failed to conduct these programs. According to the results, it is important to show that instructions regarding any type of activities are very important. As the target students are in primary school, school administrators and teachers must play their roles in giving specific explanations and specific instructions in order for the students to clearly understand the purpose of conducting the programs. School administrators must also take the initiative of implementing more programs related to health. These programs should not be seen as a burden because active involvement from students will benefit them for years to come and will help the students in their future development. As the results show that students are supportive, school administrators should see this as a positive sign to help the students.

LIMITATIONS

Certain limitations were found in this study. This study was limited to upper primary school students of standards 4, 5 and 6 and could be seen as not representing the whole population of primary school students. Furthermore, only 400 participants were chosen, and this study focused only on the state of Selangor. Moreover, the questionnaires were conducted in class under the supervision of teachers and not the representatives of the study. There is a possibility that the students provided answers that did not reflect their actual eating habits.

This study was only limited to student responses, but having responses from parents and the school administration would be more beneficial. With the additional responses of those two elements, the study could correlate the answers of the students, parents and administrators with respect to healthy eating. The data analysis was also limited to only frequencies and percentages.

Another major limitation is that all respondents may not have the similar ability to correctly recall their eating habits or some may over report their status. In questionnaire, the inclusion of grains (rice and noodles) in lunch box has been denoted as healthy. The term ‘whole grains’ would be more appropriate for further study as highly refined grains have similar effect as added sugars. Nonetheless, the study has provided important information on students’ knowledge, attitudes and practices towards healthy eating and the influence of school programs in promoting healthy eating. The results will benefit future programs in helping students at the primary school level.

CONCLUSIONS

The purpose of this study was to assess the knowledge, attitudes and practices of primary school students in healthy eating behaviors. It is concern that students have knowledge of the correct intake of foods but they still consume fried foods containing a high amount of saturated fat, cholesterol, sugar and salt. Most of the students are not having breakfast regularly that may cause serious health problem. Teachers and parents might give more attention to build up a habit of ‘breakfast in time’ among students. It was evident that school systems are lacking with regard to full implementation of their health programs. The ISHP was seen as a key element in inculcating a culture of health-conscious students. As students spend most of their time in school, fully utilizing the program is important to help students building healthy eating habit. Furthermore, students will be more committed with the help of teachers and friends. Students can also learn how to turn knowledge and theories into practice. In the students’ responses regarding their own eating habits, the majority of students (74%) considered themselves healthy. By implementing health programs properly, all of the students’ eating habits can be improved, thus creating a school with a healthy environment. The results of this study can be utilized by the Ministry of Education to look into improving the ISHP for the betterment of students learning a healthy lifestyle. The findings show that implementing more programs at the primary level to promote healthy eating can engage students in learning and practicing these healthy behaviors. Availability of healthier foods such as fruits, vegetables and whole grains in school campus can keep away students of changing the habits of having sugary drinks and fried foods. The school’s cooperation with parents in discussing healthy snacks and meals for students to eat at school and at home can create a healthy environment at school.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This project (RG300-14AFR) was funded by the Geran Penyelidikan Universiti Malaya (UMRG)–AFR (Frontier Science). The UM Research Cluster (Frontier Science) provided support in conducting this research. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Geran Penyelidikan Universiti Malaya (UMRG)–AFR (Frontier Science): RG300-14AFR.
UM Research Cluster (Frontier Science).

Competing Interests

There is no conflict of interest in any matter among authors.

Author Contributions

- Kazi Enamul Hoque conceived and designed the experiments, wrote the paper, reviewed drafts of the paper.
- Megat Ahmad Kamaluddin performed the experiments.
- Ahmad Zabidi Abdul Razak prepared figures and/or tables.
- Afiq Athari Abdul Wahid analyzed the data, contributed reagents/materials/analysis tools, wrote the paper.

Human Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

The study protocol was approved by the University of Malaya Research Ethics Committee.

Data Deposition

The following information was supplied regarding data availability:

The raw data is included in the frequency distribution and percentage tables in the manuscript.

REFERENCES

- Ahmad Husairi K. 2014.** Program Bersepadu Sekolah Sihat. Available at <https://prezi.com/qvs0ue4jes3y/program-bersepadu-sekolah-sihat-pbss/> (accessed 25 August 2016).
- Basch CE. 2010.** Healthier students are better learners: a missing link in school reforms to close the achievement gap. *Equity Matters: Research Review*. New York: Columbia University.
- Birch LL. 1999.** Development of food preferences. *Annual Review of Nutrition* **19**(1):41–62 DOI 10.1146/annurev.nutr.19.1.41.
- Brown JE, Isaacs J, Wooldridge N, Krinke B, Murtaugh M. 2007.** *Child and Preadolescent Nutrition, Nutrition Through the Life Cycle*. Third edition. Belmont: Thompson Wadsworth, 307–336.
- Cheng TSY, Tse LAH, Tak-Sun Y, Griffiths S. 2008.** Children's perceptions of parental attitude affecting breakfast skipping in primary sixth-grade students. *Journal of School Health* **78**(5):203–208 DOI 10.1111/j.1746-1561.2008.00287.x.
- de Graaf C. 2006.** Effects of snacks on energy intake: an evolutionary perspective. *Appetite* **47**(1):18–23 DOI 10.1016/j.appet.2006.02.007.
- Department of Health. 2009.** *Assessment of Dietary Pattern in Primary Schools 2008*. Hong Kong: Central Health Education Unit.

- Department of Statistics.** 2015. *Pusat Pentadbiran Kerajaan Persekutuan*. PutraJaya: Ministry of Health.
- Egner R, Oza-Frank R, Cunningham SA.** 2014. The school breakfast program: a view of the present and preparing for the future—a commentary. *Journal of School Health* **84**(7):417–420 DOI [10.1111/josh.12164](https://doi.org/10.1111/josh.12164).
- Ellulu M, Abed Y, Rahmat A, Ranneh Y, Ali F.** 2014. Epidemiology of obesity in developing countries: challenges and prevention. *Global Epidemic Obesity* **2**(1):2 DOI [10.7243/2052-5966-2-2](https://doi.org/10.7243/2052-5966-2-2).
- Gay LR.** 1996. *Educational Research: Competencies for Analysis and Application*. Prentice Hall: Florida International University.
- Giovannini M, Agostoni C, Shamir R.** 2010. Symposium overview: do we all eat breakfast and is it important? *Critical Reviews in Food Science and Nutrition* **50**(2):97–99 DOI [10.1080/10408390903467373](https://doi.org/10.1080/10408390903467373).
- Gourdet CK, Chriqui JF, Piekarz E, Dang Q, Chaloupka FJ.** 2014. Carrots and sticks: compliance provisions in state competitive food laws—examples for state and local implementation of the updated USDA standards. *Journal of School Health* **84**(7):466–471 DOI [10.1111/josh.12168](https://doi.org/10.1111/josh.12168).
- Holliday A, Batey C, Eves FF, Blannin AK.** 2014. A novel tool to predict food intake: the visual meal creator. *Appetite* **79**:68–75 DOI [10.1016/j.appet.2014.04.001](https://doi.org/10.1016/j.appet.2014.04.001).
- Imamura F, Micha R, Wu JHY, de Oliveira Otto MC, Otite FO, Abioye AI, Mozaffarian D.** 2016. Effects of saturated fat, polyunsaturated fat, monounsaturated fat, and carbohydrate on glucose-insulin homeostasis: a systematic review and meta-analysis of randomised controlled feeding trials. *PLoS Medicine* **13**(7):e1002087 DOI [10.1371/journal.pmed.1002087](https://doi.org/10.1371/journal.pmed.1002087).
- Karim NA.** 2014. Growing big and round. *Positive Parenting*. Available at <http://mypositiveparenting.org/2014/02/05/growing-big-and-round/> (accessed 10 February 2014).
- Kesten JM, Cameron N, Griffiths PL.** 2013. Assessing community readiness for overweight and obesity prevention in pre-adolescent girls: a case study. *BMC Public Health* **13**(1):1205 DOI [10.1186/1471-2458-13-1205](https://doi.org/10.1186/1471-2458-13-1205).
- Khor GL, Zalilah MS, Phan YY, Ang M, Maznah B, Norimah AK.** 2009. Perceptions of body image among Malaysian male and female adolescents. *Singapore Medical Journal* **50**(3):303–311.
- Krejcie RV, Morgan DW.** 1970. Determining sample size for research activities. *Educational and Psychological Measurement* **30**(3):607–610.
- Nagel G, Wabitsch M, Galm C, Berg S, Brandstetter S, Fritz M, Klenk J, Peter R, Prokopchuk D, Steiner R, Stroth S, Wartha O, Weiland SK, Steinacker J.** 2009. Determinants of obesity in the Ulm research on metabolism, exercise and lifestyle in children (URMEL-ICE). *European Journal of Pediatrics* **168**(10):1259–1267 DOI [10.1007/s00431-009-1016-y](https://doi.org/10.1007/s00431-009-1016-y).
- Noraziah A, Mohd Azlan A.** 2012. The food consumption and eating behaviour of Malaysian Urbanites: issues and concerns. *Malaysia Journal of Society and Space* **8**(6):157–165 DOI [10.1038/ejcn.2010.145](https://doi.org/10.1038/ejcn.2010.145).
- Pandey D, Buzgeia MH, Suneetha E, Ahmed H, Abd El Rahaman Al Gani H, Abd El Rahman Al Kadam H, Juma Elariby N.** 2013. Breakfast skipping pattern among Benghazi primary school children. *British Food Journal* **115**(6):837–849 DOI [10.1108/BFJ-Nov-2010-0196](https://doi.org/10.1108/BFJ-Nov-2010-0196).
- Pearson N, Biddle SJH, Gorely T.** 2009. Family correlates of breakfast consumption among children and adolescents: a systematic review. *Appetite* **52**(1):1–7 DOI [10.1016/j.appet.2008.08.006](https://doi.org/10.1016/j.appet.2008.08.006).

- Sandercock GRH, Voss C, Dye L. 2010.** Associations between habitual school-day breakfast consumption, body mass index, physical activity and cardiorespiratory fitness in English schoolchildren. *European Journal of Clinical Nutrition* **64(10)**:1086–1092 DOI [10.1038/ejcn.2010.145](https://doi.org/10.1038/ejcn.2010.145).
- Singleton N, Rhoads DS. 1982.** Meal and snacking patterns of students. *Journal of School Health* **52(9)**:529–534 DOI [10.1111/j.1746-1561.1982.tb04033.x](https://doi.org/10.1111/j.1746-1561.1982.tb04033.x).
- Woodruff SJ, Hanning RM. 2009.** Effect of meal environment on diet quality rating. *Canadian Journal of Dietetic Practice and Research* **70(3)**:118–124 DOI [10.3148/70.3.2009.118](https://doi.org/10.3148/70.3.2009.118).
- World Health Organization. 1998.** *Information Series on School Health: Healthy Nutrition: An Essential Element of a Health-Promoting School*. Geneva: World Health Organization.
- World Health Organization. 2013.** *Factsheets*. Geneva: WHO Media Centre.